

Application for Coverage of Belimumab for Lupus Nephritis

Adjunct treatment for class III or IV +/- class V lupus nephritis within 4 months of induction treatment and $eGFR \ge 30 \text{ mL/min/1.73m}^2$

Revised: May/25	Page 1 of 4

► PLEASE SIGN, DATE AND FAX THIS FORM TO: BC Renal (604) 875-7366

PATIEN	T INFORMATION LABEL
Name:	
Address	
Phone:	
Date of E	Birth (MM/DD/YYYY):
PHN:	

INSTRUCTIONS

- 1. Ensure the patient is registered in PROMIS under the Provincial Renal Agency and under the care of a nephrologist within a BC Renal Health Authority program.
 - a) Choose the most appropriate GN diagnosis under the available list of primary renal diagnoses.
 - b) Ensure the patient address and contact information are accurate as these are needed for medication distribution.
- 2. Complete the appropriate section below, fax this form to the BC Renal at (604) 875-7366.
 - a) Please note applications at 12 or 24 months should use the same type of form that was selected for initiation
- 3. This application will be reviewed by the BC Renal Pharmacy Services Committee; you will be contacted once approval is decided.
- 4. If approved, fax the approval letter along with the Benlysta Monarch Patient Support Program enrollment form with the Lupus Nephritis section completed to the **Monarch Patient Support Program at (1-855-788-3140)**, to inform them of BCR coverage. The Patient Support Program will then arrange intravenous infusion or teaching for subcutaneous self-administration.
- 5. All approvals are valid for 12 months from the approval date, you will need to reapply after that.

THE FOLLOWING ARE REQUIRED FOR MEDICATION APPROVAL:

GN Dia	gnosis with PROMIS codes (pick one):
	Lupus nephritis (84), class III
	Lupus nephritis (84), class III + V
	Lupus nephritis (84), class IV
	Lupus nephritis (84), class IV + V
	Lupus nephritis (84), class V
	Additional details about diagnosis, if needed
Weight	: Age:
Intrave	nous Administration
	10 mg/kg = mg IV every 2 weeks x first 3 doses, then every 4 weeks thereafter
	10 mg/kg = mg IV every 4 weeks
	Other:
Subcut	aneous Administration
	400 mg subcutaneous weekly x first 4 doses, then 200 mg weekly thereafter
	200 mg subcutaneous weekly
	Other:



Application for Coverage of Belimumab for Lupus Nephritis

Adjunct treatment for class III or IV +/- class V lupus nephritis within 4 months of induction treatment and $eGFR \ge 30 \text{ mL/min/1.73m}^2$

Revised: May/25	Page 2 of 4

PATIENT INFORMATION LABEL
Name:
Address:
Phone:
Date of Birth (MM/DD/YYYY):
PHN:

Initiation criteria

П	Adults with class III or IV	with or without class V.	lupus nephritis on a	ı kidnev biopsy	within the last 6 months:

Provide date of biopsy (MM/DD/YYYY):

If a biopsy can't be performed:

Justify why:

Justify a clinical diagnosis of class III or IV with or without class V lupus nephritis disease flare: ______

□ Started induction therapy for the treatment of lupus nephritis within the prior 4 months, and have not already entered remission, and has an eGFR \geq 30mL/min/1.73m²

• Provide details of drugs used for current induction therapy and their start dates and doses:

Provide the following laboratory values:

Labs:	Prior to the onset of the most recent flare	At the time of the onset of current induction treatment	The most recent values
creatinine			
eGFR			
proteinuria			
urinalysis			

- ☐ Has not previously failed both cyclophosphamide and mycophenolate (both with corticosteroids) as induction therapies for lupus nephritis
 - Provide details of previous induction therapy used in the past and the therapeutic response observed:

Continuation criteria at 12 months

- ☐ Has not started any new immunosuppression therapies for the treatment of lupus nephritis (other than those that were initially part of induction or maintenance therapy)
- ☐ Has reduced corticosteroids to ≤ 10 mg/day of prednisone or equivalent (other than short-term increases for non-lupus nephritis indications)
- □ Current eGFR ≥ 30mL/min/1.73m², has not started dialysis and has not received a kidney transplant
 - Provide the following laboratory values:

Labs:	Prior to the onset of the most recent flare
creatinine	
eGFR	



Application for Coverage of Belimumab for Lupus Nephritis

Adjunct treatment for class III or IV +/- class V lupus nephritis within 4 months of induction treatment and $eGFR \ge 30 \text{ mL/min/1.73m}^2$

Revised: May/25	Page 3 of 4

Continuation criteria at 24 months and every 12 months thereafter

Has not started any new immunosuppression therapies for the treatment of lupus nephritis (other than those
that were initially part of induction or maintenance therapy)

PATIENT INFORMATION LABEL

Date of Birth (MM/DD/YYYY):

Name:

Address:

Phone:

PHN:

☐ Has reduced corticosteroids to ≤ 10 mg/day of prednisone or equivalent (other than short-term increases for non-lupus nephritis indications)

An eGFR that has been stable or improved in the prior 12 months and is currently eGFR \geq 30 mL/min/1.73m², has not started dialysis and has not received a kidney transplant

• Provide the following laboratory values:

Labs:	Prior to onset of most recent flare	At time of most recent flare	12 months ago	The most recent values
creatinine				
eGFR				

•	If eGFR is declining due to reasons other than active lupus nephritis, provide a detailed explanation to
	justify ongoing treatment with belimumab:

And satisfies either 1 or 2 or 3 below:

- ☐ 1) Has continued to show clinical improvement but is not yet in remission, as evidenced by proteinuria improvement by $\geq 25\%$ in the prior 12 months and is ≥ 0.7 g/day.
 - Provide the following laboratory values:

Labs: 12 months ago		The most recent value
proteinuria		

OR

1 2) Has achieved remission but has a history of lupus nephritis flare on maximum maintenance the	
	as evidenced by proteinuria < 0.7 g/day and there is a history of previous lupus nephritis flare (biopsy
	proven class III or IV with or without class V) while on maximum maintenance dose of mycophenolate (or
	has a contraindication to mycophenolate and has had a flare on maximum dose of azathioprine).
	2)

•	Provide details of maintenance therapy, with drug dosing at time of prior flares, and details of biopsy
	dates and findings that confirmed the lupus nephritis flare:

•	If current proteinuria is ≥ 0.7 g/day, a kidney biopsy showing resolution of class III or IV active lesions
	(i.e. histologic remission) can be used instead, provide biopsy date and details:



Address:
Phone:
Date of Birth (MM/DD/YYYY):
PHN:
naximum maintenance therapy to maintain hab for the treatment of lupus nephritis and IV with or without class V lupus nephritis confirmed elimumab while on maximum maintenance dose of e if has a contraindication to mycophenolate) story, details regarding achieving remission while on aintenance therapy at the time of the lupus nephritis I the details of the kidney biopsy date and findings IAVE A COMPLICATED HISTORY OF LUPUS T QUALIFY FOR ONGOING USE OF BELIMUMAB.
hours, or if you'd like to check on the status of your lications@bcrenal.ca.
nature:

PATIENT INFORMATION LABEL

Name: